

Order Information		
Order Date	Orders must be picked up within 72 hours of notification phone call from Peel Public Health.	
Physician/Practice Name		
Address		
City	Postal Code	Telephone Number
Contact Person	Type of Practice	Number of Physician(s) and/or NP(s) in facility
Delivery or Pick-up Preference (select one)		
<input type="checkbox"/> Vaccine Delivery (Registered participants only) <input type="checkbox"/> Hurontario - 7120 Hurontario Street, Mississauga <input type="checkbox"/> Brampton - 150 Central Park Drive, Brampton (Open on TUESDAY, Thursday, and FRIDAY only)		
Please complete the following: (if you are a Long-Term Care facility ordering for your residents >65 years, please call to order vaccines)		
Type of Refrigerator: <input type="checkbox"/> Bar Style <input type="checkbox"/> Kitchen Style <input type="checkbox"/> Laboratory		No. of fridges used for vaccine storage: _____
Details on scheduled Influenza Community Clinic (if applicable):		
Estimated Date: _____		Estimated No. of doses required: _____
Influenza Vaccine Inventory - Specify number of DOSES in each field below:		
Current stock (doses):	No. of Doses required for Risk Groups:	No. of Doses required for the General Population:
**Seasonal Influenza Vaccine will be distributed based on available supply. ** Agriflu™ (ten 0.5 ml pre-filled syringes) or Vaxigrip® (5 ml multi-dose vial) depending on available supply.		
By submitting this order, I _____ verify on behalf of the practice that the fridge storing publicly funded vaccines, at the location listed above, maintains cold chain temperatures (between +2.0°C to +8.0°C), and meets MOHLTC Vaccine Storage and Handling Guidelines. I understand that we may be required to provide accurate temperature logs upon request and that temperature logs must be kept on-site for a minimum of 3 years.		
Signature: _____		Date: _____
Recommended Recipients of Influenza Vaccine (NACI; Volume 36 ACS August 2010)		
Persons at high risk of influenza-related complications	People capable of transmitting influenza to those at high risk:	
<ul style="list-style-type: none"> ▪ Adults (including pregnant women) and children with chronic health conditions. ▪ Cardiac or pulmonary disorders (including bronchopulmonary dysplasia, cystic fibrosis and asthma); ▪ diabetes mellitus and other metabolic diseases; ▪ cancer, immunodeficiency, immunosuppression (due to underlying disease and/or therapy); ▪ renal disease; ▪ anemia or hemoglobinopathy; ▪ conditions that compromise the management of respiratory secretions and are associated with an increased risk of aspiration; ▪ children and adolescents with conditions treated for long periods with acetylsalicylic acid. ▪ People ≥ 65 years of age. ▪ People of any age who are residents of long term care and other chronic care facilities. ▪ Healthy children 6 to 23 months of age. ▪ Healthy pregnant women; (risk of influenza-related hospitalization increases. with length of gestation, i.e. it is higher in the 3rd than in the 2nd trimester. 	<ul style="list-style-type: none"> ▪ Health care and other care providers in facilities and community settings who, through their activities, are capable of transmitting influenza to those at high risk of influenza complications. ▪ Household contacts (adults and children) of individuals at high risk of influenza-related complications (whether or not the individual at high risk has been immunized); <ul style="list-style-type: none"> ▪ Household contacts of individuals at high risk (see previous column); ▪ Household contacts of infants under 6 months of age (who are at high risk of complications from influenza but for whom influenza vaccine is not approved); ▪ Members of a household expecting a newborn during the influenza season. ▪ Those providing regular child care to children under 24 months of age, whether in or out of the home. ▪ Those who provide services within closed or relatively closed settings to persons at high risk (e.g. crew on a ship). 	
Special Consideration:	Other Recommended groups:	
<ul style="list-style-type: none"> ▪ Persons with morbid obesity (BMI ≥ 40) ▪ Aboriginal peoples ▪ Children 2-4 years of age 	<ul style="list-style-type: none"> ▪ People who provide essential community services ▪ People in direct contact with avian-influenza infected poultry during culling operations. 	
General Population		
All individuals aged 6 months or older, who live, work or attend school in Ontario.		
FOR PEEL ADMINISTRATIVE SUPPORT USE ONLY		
Vaccine Order Packed By: _____	Date Sent: _____	# of Bags _____
Agriflu™ lot# _____	Expiry Date: _____	
Vaxigrip® lot# _____	Expiry Date: _____	
Vaccine Management and Physician Information Team Telephone 905-791-7800 ext. 6404 Fax: 905-565-9874		